

Metro Public Health Department STD/HIV Intervention and Prevention
Total number of reported STD cases and rates[†] per 100,000 population: Nashville-Davidson County, Tennessee 1996-2005.*^{††}

Year Reported	Chlamydia		Gonorrhea		1° Syphilis		2° Syphilis		E Latent Syphilis		Congenital Syphilis		Other Stages of Syphilis	
	No. of cases	Rate per 100,000	No. of cases	Rate per 100,000	No. of cases	Rate per 100,000	No. of cases	Rate per 100,000	No. of cases	Rate per 100,000	No. of cases	Rate per 100,000 live births [§]	No. of cases	Rate per 100,000
1996	1,965	351.9	2,033	364.1	58	10.4	135	24.2	99	17.7	1	...	0	0.0
1997	1,820	322.1	2,050	362.8	61	10.8	142	25.1	173	30.6	0	0.0	36	6.4
1998	1,981	349.9	1,777	313.9	50	8.8	160	28.3	148	26.1	0	0.0	58	10.2
1999	2,202	388.2	1,785	314.7	86	15.2	164	28.9	202	35.6	0	0.0	54	9.5
2000	2,403	421.7	2,404	421.8	77	13.5	123	21.6	173	30.4	4	...	145	25.4
2001	2,086	363.6	1,625	283.2	19	3.3	57	9.9	137	23.9	5	...	162	28.2
2002	2,157	373.2	1,409	243.8	7	...	19	3.3	109	18.9	5	...	154	26.6
2003	2,653	455.5	1,492	256.2	6	...	15	2.6	25	4.3	0	0.0	149	25.6
2004	2,664	453.6	1,221	207.9	2	...	13	2.2	22	3.7	3	...	124	21.1
2005	2,819	475.8	1,116	188.4	3	...	21	3.5	21	3.5	0	0.0	99	16.7

*County level population estimates (denominator) used in crude rate calculations provided by Tennessee Department of Health Office of Health Statistics.

[†]Data source: 1996-2005 case counts from MPHD Epidemiology and Health Monitoring analysis of NETSS Surveillance Data, Tennessee Department of Health, HIV/AIDS/STD Surveillance and Data Management, received May 31, 2006.

[‡]Rates are not calculated for case counts less than 10 due to reliability of statistics produced from small case counts.

[§]Congenital Syphilis rates in infants less than 1 year old.

Table prepared by Jeselyn Rhodes, Division of Epidemiology and Health Monitoring